Abbreviations

3TC	Lamivudine
ABC	Abacavir
AGL	Adherence Guideline
AHD	Advanced HIV disease
ALT	Alanine transaminase
am	In the morning
ANC	Antenatal Care
APC	Adult Primary Care
ART	Antiretroviral therapy
ARV	Antiretroviral
ATV/r	Atazanavir/ritonavir
AZT	Zidovudine
bd	Twice daily
BMI	Body mass index
CM	Cryptococcal meningitis
CNS	Central nervous system
CPT	Cotrimoxazole preventive therapy
CrAg	Cryptococcal Antigen
CVS	Cardiovascular
DILI	Drug-induced liver injury
DMOC	Differentiated Models of Care
DR	Drug-resistant
DS	Drug-sensitive
DT	Dispersible tablet
DTG	Dolutegravir
EAC	Enhanced Adherence Counselling
eGFR	Estimated glomerular filtration rate
EFV	Efavirenz
EX-PUP	External pick-up point
FAC-PUP	Facility pick-up point
FC	Film coated
FDC	Fixed-dose combination
GIT	Gastrointestinal tract;
Hb	Haemoglobin
HBsAg	Hepatitis B surface antigen
HBV	Hepatitis B virus
InSTI	Integrase strand transfer inhibitor
IRIS	Immune reconstitution inflammatory syndrome
IUCD	Intrauterine contraceptive device
LP	Lumbar puncture
LPV/r	Lopinavir/ritonavir
MMD	Multi-month Dispensing
MUAC	Mid-upper arm circumference
NA	Not applicable
NCDs	Non-communicable diseases
nocte	at night
NNRTI	Non-nucleoside reverse transcriptase inhibitor
NRTI	Nucleoside reverse transcriptase inhibitor
NTDs	Neural tube defects
NVP	Nevirapine
od	Once daily
OI	Opportunistic infection
PBFW	Pregnant and breastfeeding women
PCR	Polymerase chain reaction test for HIV
PHC EML	Primary Health Care Essential Medicines List
PI	Protease inhibitor
PJP	Pneumocyctis jerovecii pneumonia
PLHIV	People living with HIV
pm	in the evening
RPCs	Repeat prescription collection strategies
RT	Resistance test
sCR	Serum creatinine

std	Standard
STIs	Sexually transmitted infections
TB	Tuberculosis
TDF	Tenofovir disoproxil fumarate
TEE	Tenofovir + emtricitabine + efavirenz
TLD	Tenofovir + lamivudine + dolutegravir
TPT	TB preventive treatment
VL	Viral load
VT	Vertical transmission
WHO	World Health Organisation