
Appendix A:



Mini-Cog©

Instructions for Administration & Scoring

ID: _____ Date: _____

Step 1: Three Word Registration

Look directly at person and say, "Please listen carefully. I am going to say three words that I want you to repeat back to me now and try to remember. The words are [select a list of words from the versions below]. Please say them for me now." If the person is unable to repeat the words after three attempts, move on to Step 2 (clock drawing).

The following and other word lists have been used in one or more clinical studies.[1-3] For repeated administrations, use of an alternative word list is recommended.

Version 1	Version 2	Version 3	Version 4	Version 5	Version 6
Banana	Leader	Village	River	Captain	Daughter
Sunrise	Season	Kitchen	Nation	Garden	Heaven
Chair	Table	Baby	Finger	Picture	Mountain

Step 2: Clock Drawing

Say: "Next, I want you to draw a clock for me. First, put in all of the numbers where they go." When that is completed, say: "Now, set the hands to 10 past 11."

Use preprinted circle (see next page) for this exercise. Repeat instructions as needed as this is not a memory test. Move to Step 3 if the clock is not complete within three minutes.

Step 3: Three Word Recall

Ask the person to recall the three words you stated in Step 1. Say: "What were the three words I asked you to remember?" Record the word list version number and the person's answers below.

Word List Version: _____ Person's Answers: _____ _____ _____

Scoring

Word Recall: _____ (0-3 points)	1 point for each word spontaneously recalled without cueing.
Clock Draw: _____ (0 or 2 points)	Normal clock = 2 points. A normal clock has all numbers placed in the correct sequence and approximately correct position (e.g., 12, 3, 6 and 9 are in anchor positions) with no missing or duplicate numbers. Hands are pointing to the 11 and 2 (11:10). Hand length is not scored. Inability or refusal to draw a clock (abnormal) = 0 points.
Total Score: _____ (0-5 points)	Total score = Word Recall score + Clock Draw score. A cut point of <3 on the Mini-Cog™ has been validated for dementia screening, but many individuals with clinically meaningful cognitive impairment will score higher. When greater sensitivity is desired, a cut point of <4 is recommended as it may indicate a need for further evaluation of cognitive status.

Mini-Cog © S. Borson. All rights reserved. Reprinted with permission of the author solely for clinical and educational purposes. May not be modified or used for commercial, marketing, or research purposes without permission of the author (soob@uw.edu).
v. 01.19.16


Clock Drawing

ID:_____ Date:_____

References

1. Borson S, Scanlan JM, Chen PJ et al. The Mini-Cog as a screen for dementia: Validation in a population based sample. J Am Geriatr Soc 2003;51:1451–1454.
2. Borson S, Scanlan JM, Watanabe J et al. Improving identification of cognitive impairment in primary care. Int J Geriatr Psychiatry 2006;21: 349–355.
3. Lessig M, Scanlan J et al. Time that tells: Critical clock-drawing errors for dementia screening. Int Psychogeriatr. 2008 June; 20(3): 459–470.
4. Tsoi K, Chan J et al. Cognitive tests to detect dementia: A systematic review and meta-analysis. JAMA Intern Med. 2015; E1-E9.
5. McCarten J, Anderson P et al. Screening for cognitive impairment in an elderly veteran population: Acceptability and results using different versions of the Mini-Cog. J Am Geriatr Soc 2011; 59: 309-213.
6. McCarten J, Anderson P et al. Finding dementia in primary care: The results of a clinical demonstration project. J Am Geriatr Soc 2012; 60: 210-217.
7. Scanlan J & Borson S. The Mini-Cog: Receiver operating characteristics with the expert and naive raters. Int J Geriatr Psychiatry 2001; 16: 216-222.

Mini-Cog © S. Borson. All rights reserved. Reprinted with permission of the author solely for clinical and educational purposes. May not be modified or used for commercial, marketing, or research purposes without permission of the author (soob@uw.edu).
v. 01.19.16



CLINICAL FRAILTY SCALE

1 VERY FIT — People who are robust, active, energetic and motivated. They tend to exercise regularly and are among the fittest for their age.

2 FIT — People who have **no active disease symptoms** but are less fit than category 1. Often, they exercise or are very **active occasionally**, e.g., seasonally.

3 MANAGING WELL — People whose **medical problems are well controlled**, even if occasionally symptomatic, but often are **not regularly active** beyond routine walking.

4 LIVING WITH VERY MILD FRAILTY — Previously "vulnerable," this category marks early transition from complete independence. While **not dependent** on others for daily help, often **symptoms limit activities**. A common complaint is being "slowed up" and/or being tired during the day.

5 LIVING WITH MILD FRAILTY — People who often have **more evident slowing**, and need help with **high order instrumental activities of daily living** (finances, transportation, heavy housework). Typically, mild frailty progressively impairs shopping and walking outside alone, meal preparation, medications and begins to restrict light housework.

6 LIVING WITH MODERATE FRAILTY — People who need help with **all outside activities** and with **keeping house**. Inside, they often have problems with stairs and need **help with bathing** and might need minimal assistance (cuing, standby) with dressing.

7 LIVING WITH SEVERE FRAILTY — **Completely dependent for personal care**, from whatever cause (physical or cognitive). Even so, they seem stable and not at high risk of dying (within ~6 months).

8 LIVING WITH VERY SEVERE FRAILTY — Completely dependent for personal care and approaching end of life. Typically, they could not recover even from a minor illness.

9 TERMINALLY ILL — Approaching the end of life. This category applies to people with a **life expectancy <6 months**, who are **not otherwise living with severe frailty.** (Many terminally ill people can still exercise until very close to death.)

SCORING FRAILTY IN PEOPLE WITH DEMENTIA

The degree of frailty generally corresponds to the degree of dementia. Common **symptoms in mild dementia** include forgetting the details of a recent event, though still remembering the event itself, repeating the same question/story and social withdrawal.

In **moderate dementia**, recent memory is very impaired, even though they seemingly can remember their past life events well. They can do personal care with prompting.

In **severe dementia**, they cannot do personal care without help.

In **very severe dementia** they are often bedfast. Many are virtually mute.

DALHOUSIE UNIVERSITY
www.geriatricmedicineresearch.ca

Clinical Frailty Scale ©2005–2020 Rockwood, Version 2.0 (EN). All rights reserved. For permission: www.geriatricmedicineresearch.ca
Rockwood K et al. A global clinical measure of fitness and frailty in elderly people. CMAJ 2005;173:489–495.

4AT

Assessment test for delirium & cognitive impairment

(label)

Patient name:

Date of birth:

Patient number:

Date: **Time:**

Tester:

	CIRCLE

[1] ALERTNESS

This includes patients who may be markedly drowsy (eg. difficult to rouse and/or obviously sleepy during assessment) or agitated/hyperactive. Observe the patient. If asleep, attempt to wake with speech or gentle touch on shoulder. Ask the patient to state their name and address to assist rating.

Normal (fully alert, but not agitated, throughout assessment)	**0**
Mild sleepiness for <10 seconds after waking, then normal	**0**
Clearly abnormal	**4**

[2] AMT4

Age, date of birth, place (name of the hospital or building), current year.

No mistakes	**0**
1 mistake	**1**
2 or more mistakes/untestable	**2**

[3] ATTENTION

Ask the patient: "Please tell me the months of the year in backwards order, starting at December." To assist initial understanding one prompt of "what is the month before December?" is permitted.

Months of the year backwards	Achieves 7 months or more correctly	**0**
	Starts but scores <7 months / refuses to start	**1**
	Untestable (cannot start because unwell, drowsy, inattentive)	**2**

[4] ACUTE CHANGE OR FLUCTUATING COURSE

Evidence of significant change or fluctuation in: alertness, cognition, other mental function (eg. paranoia, hallucinations) arising over the last 2 weeks and still evident in last 24hrs

No	**0**
Yes	**4**

4 or above: possible delirium +/- cognitive impairment
1-3: possible cognitive impairment
0: delirium or severe cognitive impairment unlikely (but delirium still possible if [4] information incomplete)

4AT SCORE ☐

GUIDANCE NOTES Version 1.2. Information and download: **www.the4AT.com**

The 4AT is a screening instrument designed for rapid initial assessment of delirium and cognitive impairment. A score of 4 or more *suggests* delirium but is not diagnostic: more detailed assessment of mental status may be required to reach a diagnosis. A score of 1-3 suggests cognitive impairment and more detailed cognitive testing and informant history-taking are required. A score of 0 does not definitively exclude delirium or cognitive impairment: more detailed testing may be required depending on the clinical context. Items 1-3 are rated *solely on observation of the patient at the time of assessment.* Item 4 requires information from one or more source(s), eg. your own knowledge of the patient, other staff who know the patient (eg. ward nurses), GP letter, case notes, carers. The tester should take account of communication difficulties (hearing impairment, dysphasia, lack of common language) when carrying out the test and interpreting the score.

Alertness: Altered level of alertness is very likely to be delirium in general hospital settings. If the patient shows significant altered alertness during the bedside assessment, score 4 for this item. **AMT4 (Abbreviated Mental Test - 4):** This score can be extracted from items in the AMT10 if the latter is done immediately before. **Acute Change or Fluctuating Course:** Fluctuation can occur without delirium in some cases of dementia, but marked fluctuation usually indicates delirium. To help elicit any hallucinations and/or paranoid thoughts ask the patient questions such as, "Are you concerned about anything going on here?"; "Do you feel frightened by anything or anyone?"; "Have you been seeing or hearing anything unusual?"

© 2011-2014 MacLullich, Ryan, Cash.